

WHEAT DISEASE CONTROL

Alfredo Martinez-Espinoza, Extension Plant Pathologist

DISEASE	CHEMICAL AND FORMULATION	RATE PER ACRE	REMARKS AND PRECAUTIONS
Stagonospora Leaf and Glume Blotch, Leaf Rust, Stripe Rust, Powdery Mildew, Tan Spot	Azoxystrobin Quadris	6.2-10.8 oz	Apply after Feekes 6 but not later than Feekes 10.5. Do not harvest treated wheat for forage. A crop oil concentrate adjuvant may be added at 1.0% v/v to optimize efficacy.
	Fluoxapyroxad + Pyraclostrobin Priaxor	4-8 fl. oz	Apply no later than the beginning of flowering (Feekes 10.5 Zadok's). Maximum number of applications per season =2.
	Fluoxastrobin Evito	2-4 fl. oz	For optimum results, begin applications preventatively and continue on a 14 to 21 day interval. Do not make more than two sequential applications. Apply prior to disease development from Feekes 5 (Zadok's 31) up to late head emergence at Feekes 10.5 (Zadok's 59).
	Metconazole Caramba	10-14 oz	Maximum number of applications per season=2; Minimum time from application to harvest=30 days.
	Propiconazole Tilt Propimax	4 oz	Tilt can be applied until heading stage (Feekes 10.5). Do not apply Tilt after this growth stage to avoid possible illegal residues.
	Propiconazole-Azoxystrobin Quilt, QuiltXcel	7-14 oz	Applications may be made no closer than a 14 day interval. Quilt and QuiltXcel can be applied up to Feekes growth stage 10.5. QuiltXcel has a higher rate of azoxystrobin. Low rates of Quilt and QuiltXcel are used for spring suppression of early season diseases. 10.5 fl. oz. and above are used for flag leaf protection and maximizing yield potential.
	Propiconazole-Trifloxystrobin Stratego	10 oz	Do not apply more than 2 applications of Stratego per season. Do not apply after Feekes 10.5.
	Prothioconazole Proline	4.3-5 fl oz	For optimum disease control, the lowest labeled rate of a spray surfactant should be tank mixed with Proline. Up to two applications of Proline can made per year.
	Prothioconazole + Tebuconazole Prosaro	6.5-8.2 fl oz	Begin applications preventively when conditions are favorable for disease development. For optimum disease control, the lowest labeled rate of a spray surfactant should be tank mixed with Prosaro.
	Prothioconazole + Trifloxystrobin Stratego YLD	4 fl oz	Begin applications preventively when conditions are favorable for disease development. Do not apply more than 2 applications per season. Do not apply after Feekes growth stage 10.5. Do not apply within 35 days of harvest.
	Pyraclostrobin Headline	6-9 oz	Apply no later than Feekes 10.5.
	Pyraclostrobin + Metconazole Twinline	7-9 fl oz	Do not apply more than 2 applications per season. Do not apply after Feekes 10.5.
	Tebuconazole Folicur	4 fl oz	Folicur is not longer manufactured (2009). No end-user restrictions for disease control. Use until supply runs out. Not labeled for Powdery mildew control.
	Tebuconazole + Trifloxystrobin Absolute	3-5 fl oz	Begin applications preventively when conditions are favorable for disease development. For optimum disease control apply 5 fl. oz. at flag leaf stage (Feekes 8-9). For early season suppression of Tan Spot, Leaf Blight and Powdery Mildew, apply at 3-4 oz. Do not apply more than 5 fl. oz. per season. Do not apply after Feekes growth stage 10.5.2. Do not apply within 35 days of harvest. Do not use with adjuvants.